

# International Cooperators'



# **Procedures for Tomato Variety Field Trials**

Peter Hanson, Li-ju Lin, Gregory C. Luther, Wen-shi Tsai, R. Srinivasan, Chien-hua Chen, Chih-hung Lin, Zong-ming Sheu and Shu-fen Lu

### Introduction

The procedures described here allow comparison of the data collected in different test environments (locations, years, and seasons) by researchers participating in WorldVeg multienvironment tomato variety trials. They could also be useful for other researchers interested in testing tomato varieties under local conditions.

#### Choice of land

Select a well-drained area with fairly uniform fertility and slope.

#### Number of entries

The suggested number of entries is from 5 to 20, which should include one or two locally popular varieties at each location (Table 1).

#### Experimental design

A randomized complete block design (RCBD) with three replications is recommended (Fig. 1). Each field trial has border rows on four sides.

#### Size of plot

Row length and plant spacing normally used in local production practices are recommended.

The minimum number of plants per plot is 12 (1-row planting for large entries, data is collected from the 10 inner plants). At WorldVeg, each entry is grown on a 2-row, 4.8 m long and 1 m wide plot with furrows (ditches) 50 cm wide on each side. The distance between rows is 60 cm. Plant spacing within rows is 40 cm. Thus, each row accommodates 12 plants and a total of 24 plants per plot. Data is collected from the 20 inner plants. Any changes in plot dimensions should be reflected in the data sheet.

#### **Cultural practices**

For recommended cultural and pest management practices, please refer to:

- Suggested Cultural Practices for Tomato
- Pruning and Staking Tomatoes
- Safer Tomato Production Techniques

in **HARVEST**, the WorldVeg document and data archive: <u>https://worldveg.tind.io/</u>

Search: PRODUCTION GUIDES

Table	1.	Sampl	e p	lanting	plan.
-------	----	-------	-----	---------	-------

	Replication									
Entry code	Ι	II	III							
	Plots 1-8	Plots 9-16	Plots 17-24							
A	8	12	17							
В	7	16	22							
С	5	10	20							
D	4	15	21							
Е	2	11	19							
F	3	13	18							
G	1	14	24							
H*	6	9	23							

\*Local check variety



Figure. 1. Sample field layout (the border area can have one or two rows of tomatoes).

## Harvesting

For fresh market tomato, start harvesting at breaker stage (less than 10% surface pink or red). For cherry tomato, harvest the whole fruit cluster when 80-90% turns red. For processing tomato, harvest red ripe fruit. At WorldVeg, determinate tomato plants are generally harvested three times and indeterminate types four or more times. Record harvest dates and times (Table 2).

## Data to collect

Researchers should keep a record of the basic characteristics of the trial site and the management practices employed when conducting a variety trial (Tables 2 & 3). This information can be useful for explaining varietal performance in different environments. Plant characteristics and reactions to biotic stresses, yield and its components to be collected for each plot are as follows:

#### 1. Days to 50% flowering:

Number of days after transplanting when 50% of the plants in a plot have open flowers. Check plots three times a week (Table 4).

#### 2. Growth habit:

(1) *determinate:* short and bushy, produces two leaves between flower clusters and about five clusters per branch; (2) *indeterminate:* tall, produces three leaves between flower clusters and more than six clusters per branch; (3) *semi-determinate:* taller than determinate types, but not as tall as indeterminate types (Table 4).

#### 3. Biotic stress rating:

Entries are evaluated every 1-2 weeks when pest pressure (damage) is most serious. Check Figures 2 to 4 for the rating scales of early blight, late blight and tomato yellow leaf curl disease (TYLCD); and Figures 5 to 7 for wilt symptoms of fusarium wilt, bacterial wilt and southern blight to help you score and record the severity of diseases (Table 4). For insect damage, count and weigh the number of fruits damaged by tomato fruit borer (Tables 4 & 5).

#### 4. Number of plants harvested:

Count the plants harvested from the 2-row plot. This will indicate population density and help explain low yields in plots with poor stands (Table 5).

#### 5. Number of fruits and fruit yield:

Separate the marketable (worth selling) from nonmarketable fruits (with defects such as cracking, blossom end rot, graywall, blotchy ripening, puffiness, sunscald, catface, insect damaged fruits, etc.) after harvesting (Figure 8). Record the number and weight (kg/plot) of marketable and nonmarketable fruits. Repeat the process every time until harvesting is done. The total marketable yield is obtained by adding the yields of individual harvests (Table 5).

The yield per plot (kg/plot) can be converted into tonnes per hectare with the following formula:

Yield (t/ha) = 
$$\frac{\text{plot yield (kg) / 1,000 (kg/t)}}{\text{harvested area (m2) / 10,000 (m2/ha)}}$$

*Example:* plot yield: 30 kg

harvested area: 20 m<sup>2</sup>

$$Yield = \frac{30 (kg) / 1,000 (kg/t)}{20 (m^2) / 10,000 (m^2/ha)} = 15 t/ha$$

#### 6. Fruit weight:

Average fruit weight (grams) can be calculated from 20 randomly selected marketable fruit per plot (Table 5).

*Example:* Weight of 20 marketable fruits = 1,250 g

Average fruit weight =  $\frac{1,250}{20}$  = 62.5 g

#### 7. Remarks:

Any other interesting observations not recorded elsewhere that could help explain the outcome of the trial.

TOMATO VARIETY FIELD TRIALS: TEST LOCATION AND CRO	P MANAGEMENT DATA SHEET (1)					
Country	FIELD PLOT DATA					
State / province / department	Plot width (m)					
District / town / city	Row length (m)					
Farm or experiment station	No. of rows / plot					
Institution	No. of plants / row					
Cooperator (s)/ data taker (s)	Spacing between rows (cm)					
E-mail:	Plant spacing within rows (cm)					
LATITUDE degrees minutes N or S LONGITUDE degrees minutes	E or W ALTITUDE above sea level					
SOIL Classification Previo	bus crop					
surface texture surface p	H drainage condition					
sandy unknown						
loam 7.1 - 8	very good					
clay loam 5.6 - 7	average					
silty clay 4 - 5.5	poor					
clay< 4	very poor					
other or						
If other, specify actual value						
PLANTING SCHEDULE     day     month     year       date sown     Image: Im	day month year					
HARVEST day month year day mo	onth year Number of times					
start date	harvested					
FERTILIZER APPLIED? 🗌 Yes 🗌 No Specify unit of fertilizer ap	oplied if different from kg/ha					
applied     day     month     year     kg/ha     %N       1st date            2nd date            3rd date	%P205       %K20       Other element(s)					
IRRIGATION ?	quency					
methods	frequency					
Eurrow twice a month						
Sprinkler monthly						
Other other						
If other, specify						
OTHER PRACTICES						

#### **Table 3.** Data collection sheet for test location and crop management (2)

TOMATO VARIETY FIELD TRIALS: TEST LOCATION AND CROP MANAGEMENT DATA SHEET (2)											
Country	FIELD PLOT DATA										
State / province / department	Plot width (m)										
District / town / city	Row length (m)										
Farm or experiment station	No. of rows / plot										
Institution	No. of plants / row										
Cooperator (s)/ data taker (s)	Spacing between rows (cm)										
E-mail:	Plant spacing within rows (cm)										
foliar disease     root disease     insect or mite damage     rat or bird damage       none	herbicide damage weed problem										
MAJOR DISEASE OBSERVED (OR SYMPTOMS)											
IF A INSECT OR MITE PROBLEM IS MODERATE OR SEVERE, PLE	EASE SPECIFY:										
MAJOR INSECT OBSERVED											
CONTROL MEASURES AND DATE(S) APPLIED											
IF A WEED PROBLEM IS MODERATE OR SEVERE, PLEASE SPEC	CIFY:										
MAJOR SPECIES, CONTROL MEASURES AND DATE(S) APPLIED											
CHEMICALS APPLIED ?  Yes No											
HERBICIDE  Yes No If Yes, specify product (s)											
FUNGICIDE 🗌 Yes 🗌 No 🛛 If Yes, specify product (s)											
INSECTICIDE  Yes No If Yes, specify product (s)											
OTHERS 🗌 Yes 🗌 No 🛛 If Yes, specify product (s)											
DATES APPLIED (DD/MM/YY) Herbicide Fungicide	Insecticide Others										
1 <sup>st</sup> spray											
2 <sup>nd</sup> spray											
3 <sup>rd</sup> spray											
CLIMATE DATA DURING TRIAL   Rainy season  Dry seasor	n										
Average min. temp. $\Box \Box^{\circ}C$ Average max. temp. $\Box \Box^{\circ}C$	Total rainfall										
Remarks about deviations from normal											

Plot	Rep	ep Entry code	Days to 50% flowering	Growth habit <sup>1</sup>		Insect damage					
no.	<b>'</b>				EB <sup>2</sup>	LB <sup>2</sup>	TYLCD <sup>3</sup>	BW <sup>4</sup>	$FW^4$	SB <sup>4</sup>	TFB⁵
1	1	G									
2	1	E									
3	1	F									
4	1	D									
5	1	С									
6	1	н									
7	1	В									
8	1	А									
9	2	Н									
10	2	С									
11	2	E									
12	2	A									
13	2	F									
14	2	G									
15	2	D									
16	2	В									
17	3	А									
18	3	F									
19	3	E									
20	3	С									
21	3	D									
22	3	В									
23	3	н									
24	3	G									

	Table 4. Data collection sheet for	plant characteristics and	reactions to biotic stresses.
--	------------------------------------	---------------------------	-------------------------------

<sup>1</sup>D: determinate type; ID: indeterminate type; SD: semi-determinate type

 $^{2}$ EB (= early blight) and LB (= late blight): rate the plants at one of three levels, 0 = healthy, 1 = slight, 2 = severe  $^{3}$ TYLCD (= tomato yellow leaf curl disease): rate the plants at one of four levels, 0 = healthy, 1 = slight, 2 = moderate, 3 = severe

- <sup>4</sup> BW (= bacterial wilt), FW (= fusarium wilt) and SB (= southern blight): record number of wilted plants
- <sup>5</sup> TFB (= tomato fruit borer): record number of TFB damaged fruits

			No. of	Average	Fruit yield (kg/plot)										Total		
Plot no. Rep		Entry code	plants	fruit weight	1 <sup>st</sup> harvest( )		2 <sup>nd</sup> harvest ( )			3 <sup>rd</sup> harvest ( )			4 <sup>th</sup> harvest ( )			fruit weight	
				(g)	M <sup>1</sup>	NM <sup>2</sup>	TFB <sup>3</sup>	M <sup>1</sup>	NM <sup>2</sup>	TFB <sup>3</sup>	M <sup>1</sup>	NM <sup>2</sup>	TFB <sup>3</sup>	M <sup>1</sup>	NM <sup>2</sup>	TFB <sup>3</sup>	(kg)
1	1	G															
2	1	E															
3	1	F															
4	1	D															
5	1	С															
6	1	н															
7	1	В															
8	1	А															
9	2	н															
10	2	с															
11	2	E															
12	2	А															
13	2	F															
14	2	G															
15	2	D															
16	2	в															
17	3	А															
18	3	F															
19	3	E															
20	3	С															
21	3	D															
22	3	В															
23	3	н															
24	3	G			L									L			

#### **Table 5.** Data sheet to track yield and yield components.

( ) indicate the date of harvest. Add more columns if there are more than 4 harvests. <sup>1</sup> M: marketable fruits

<sup>2</sup> NM: nonmarketable fruits

<sup>3</sup> TFB: tomato fruit borer damaged fruits



Figure 2. Early blight rating scale: 0 = no symptoms, 1 = dark circular spots start on the old leaves, 2 = leaves dry and falling off.



Figure 3. Late blight rating scale: 0 = no symptoms, 1 = irregular dark, water-soaked spots develop on leaves and the undersides of lesions may be covered by a white fuzzy growth, 2 = brown to black lesions appear on stems and shiny, dark or olive-colored lesions develop on fruits.



Figure 4. Tomato yellow leaf curl disease rating scale: 0 = no symptoms, 1 = curling of upper leaves, 2 = curling, blistering and yellowing of leaves, 3 = stunting and distortion.



**Figure 5.** Symptoms of fusarium wilt: Yellowing begins on lower leaves and eventually leads to leaf drop and plant wilt.

**Figure 6.** Typical symptoms of bacterial wilt first appear as drooping of a few young leaves. A sudden complete wilt soon follows. Infected plants display wilting but not yellowing leaves. Most of the time, leaves are still green when the plants wilt.



**Figure 7.** Symptoms of southern blight: White fungal growth is produced on the stem at the soil line and mustard seed-sized, round, tan to dark brown structures appear on the white fungal growth, leading to a rapid wilting of the entire plant.



#### Figure 8. Nonmarketable tomato fruits.

- 1. Cracking
- 2. Blossom end rot
- 3. Graywall
- 4. Catface
- 5. Puffiness
- 6. Sunscald
- 7. Blotchy ripening
- 8. Damage by tomato fruit borer

This guide and Excel format data spreadsheets are available online in HARVEST, the WorldVeg document and data archive:

#### https://worldveg.tind.io

Search: PRODUCTION GUIDES, DATASHEETS

World Vegetable Center Headquarters P.O. Box 42, Shanhua, Tainan 74199 Taiwan **T** +886 6 5837801 **F** +886 6 5830009



## worldveg.org